


MOLECULAR NEUROSCIENCE



Patricia Revest
& Alan Longstaff

Contents

Abbreviations	ix
Preface	xv
1 Introduction	1
2 Receptor cloning	5
Introduction	5
The amino acid sequence of a protein can be deduced by sequencing its DNA	5
Restriction enzymes can be used to cut DNA	6
Separation of DNA fragments by size	6
DNA molecules can be joined using ligases	8
rDNA is introduced into bacterial cells for cloning	10
Vectors are constructed using genetic engineering techniques	10
Several methods have been devised to sequence nucleic acids	13
A strategy for cloning a desired sequence	15
The nicotinic cholinergic receptor was sequenced by probing a cDNA library	21
Box 2.1. Labeling of oligonucleotide probes	28
3 Molecular anatomy of the nervous system	31
Introduction	31
Many neurotransmitters can be located by histochemistry	31
ELISA tests can detect a wide range of molecules	33
Some neurotransmitters can be located by specific uptake mechanisms	33
Receptors can be counted using receptor ligand-binding assays	34
<i>In situ</i> hybridization can determine patterns of gene expression	35
Different kinds of blotting can identify DNA, RNA and proteins	36
Differential hybridization and subtracted cDNA libraries can measure differences in gene expression	37
The polymerase chain reaction can produce large amounts of a specific DNA	38
Single-cell PCR	41
mRNA differential display uses PCR to study changes in gene expression	41
A large number of genes expressed in human brain have been identified	43

4 Voltage-gated ion channels	
All excitable cells contain voltage-sensitive ion channels	
Sodium channels were the first to be identified	
Structure and function can be related at the molecular level	
Voltage-gated calcium channels are responsible for diverse functions	
Calcium channels have been cloned	
How similar are sodium and calcium channels?	
There is a large diversity of voltage-gated potassium channels	
Genes for potassium channels have been identified in naturally occurring mutant flies	
Not all K ⁺ channels belong to the same superfamily	
Box 4.1. Voltage clamping	
Box 4.2. Patch clamping	
Box 4.3. Site-directed mutagenesis	
Box 4.4. Chromosome walking	
5 Ionotropic receptors	
Introduction	
Nicotinic acetylcholine receptors have been studied extensively	
The M2 segments contribute to the ion pore	
GABA is the major CNS inhibitory transmitter	
GABA _A receptors bind many different ligands	
GABA _A receptor heterogeneity is evident from pharmacology	
GABA _c receptors	
Multiple glutamate receptors exist	
Separate AMPA and kainate receptors can be distinguished	
NMDA receptors	
Properties of NMDARs were studied by coexpression	
6 Metabotropic receptors and signal transduction mechanisms	
Introduction	
G protein-linked receptors activate a cascade of protein interactions	
G protein-linked receptors form a superfamily of receptors	
G protein-linked receptors share a common structure	
There are many distinct G proteins	
The actions of G proteins can be studied using a variety of techniques	
G protein α -subunits are linked to a range of effectors	
G protein $\beta\gamma$ -subunits can alter the activity of α -subunits	
Many ion channels are modulated by G proteins	
Many growth factor receptors activate an intrinsic tyrosine kinase	
Cyclic GMP levels are controlled by two separate pathways	
Mutations in 7TM receptors underlie a variety of diseases	
Dopamine receptors have been implicated in schizophrenia	
Box 6.1. Antisense knockouts	
7 Neurotransmitter release	
Introduction	
Neurotransmitter release at the neuromuscular junction is quantal	

Neurotransmitter release at central synapses may consist of a single quantum	129
Neurotransmitter release is calcium dependent	129
Which calcium channels are responsible for neurotransmitter release?	131
A large number of proteins have been identified in nerve terminals	133
Neurotransmitter release involves a number of calcium-dependent steps	134
Vesicle recruitment may involve synapsin	134
SNAPs and SNAREs are involved in docking and then priming vesicles for release	136
Synaptophysin and physophilin may form the fusion pore	138
Synaptotagmin may be the calcium sensor involved in the final exocytotic event	139
Rab3A is involved in recycling of neurotransmitter vesicles	140
Neurexins are a target for neurotoxins	141
Recycling of synaptic vesicles may involve dynamin	142
Refilling of vesicles with neurotransmitter occurs via an antiport mechanism	143
Termination of neurotransmitter action is due to diffusion and binding	144
Lambert–Eaton myasthenic syndrome is a failure of neurotransmitter release	146
Box 7.1. Imaging calcium in cells	147
Box 7.2. Production of transgenic animals	148
8 Mechanisms of plasticity	151
The hippocampus is required for episodic memory	151
Long-term potentiation is an activity-dependent increase in synaptic strength	153
Both pre- and postsynaptic events appear to be involved in the maintenance of NMDAR-dependent LTP	156
Protein kinases are activated during LTP	157
Invertebrates currently provide the most complete evidence for nuclear signaling via cAMP in plasticity	160
A metabotropic receptor switch is required for LTP	165
Presynaptic mechanisms for induction of LTP	166
Long-term depression is a synaptic weakening	173
Voltage-dependent channels are implicated in plasticity	178
LTP may be a molecular model for some types of learning	179
Epilepsy is a chronic hyperexcitable state	182
Box 8.1. Electrophoretic mobility shift assay (EMSA)	188
Box 8.2. Floxed receptor knockouts	189
9 Molecular mechanisms in neurodegenerative disease	191
Introduction	191
Huntington's disease causes a specific pattern of neuronal loss	191
Excitotoxicity involves inappropriate activation of glutamate receptors	194
Parkinson's disease also causes degeneration of specific neurons	198
Alzheimer's disease causes severe dementia	202
AD may be caused by a number of different genes	204

The amyloid hypothesis depends on the abnormal deposition of β A
Aggregation of tau proteins underlies the formation of tangles
ApoE interacts with both β A and tau in an isoform-specific manner
Other hypotheses have been suggested for the cause of AD
New therapies may be based on understanding the molecular basis of AD
Common paths to oxidative stress and free radical damage in
neurodegenerative disease

Box 9.1. Linkage analysis of DNA using RFLPs

Appendices

Appendix 1. Genetic code

Appendix 2. Single-letter code and three-letter abbreviations for
amino acids

Appendix 3. Some second messenger-activated enzymes

Glossary

Further reading

Index